


Natural Der f 2 Molecular Reference Standard

Product Code: MRS-NDF2

The Natural Der f 2 MRS is intended to serve as reference standard to determine the Der f 2 content of allergen preparations from house dust mite (*Dermatophagoides farinae*) by immunoassay.

Allergen:	Natural Der f 2 (<i>Dermatophagoides farinae</i> allergen 2)
Lot No:	xxxxx
Source:	<i>D. pteronyssinus</i> culture
Mol. Wt:	14 kD
Purification:	From spent mite culture by multi-step affinity chromatography.
Composition:	10µg natural Der f 2, determined by Amino Acid Analysis, freeze dried in sealed glass vial.
SDS-PAGE:	See inset. Silver-stained SDS-PAGE under non-reducing conditions shows a single band at 14 kD.
ELISA:	Immunoreactive in Der f 2 specific ELISA. No trace contamination with Der f 1 was detected by ELISA.
Purity:	>95% purity by in-solution LC-MS/MS after tryptic digest.
Formulation:	Prior to lyophilization, natural Der f 2 was adjusted to 50 mM volatile ammonium bicarbonate with 3% trehalose.
Storage:	Store at -20°C



nDer f 2

Natural Der f 2 Molecular Reference Standard

For research and commercial use *in vitro*: not for human *in vivo* or therapeutic use.

An InBio® product. Made in the USA

www.inbio.com

700 Harris Street
Charlottesville, VA
22903
United States

(434) 984-2304
mail@inbio.com

Vision Court
Caxton Place
Cardiff, Wales CF23 8HA
United Kingdom

+44 (0) 2921 674 640
Info@inbio.com

Product Code: MRS-NDF2-1

Reconstitution:

- Allow vial to reach room temperature before use.
- Tap vial gently to collect all material at the bottom.
- Using a sterile syringe reconstitute the MRS to desired concentration by injecting a suitable volume of a buffer of choice (e.g. PBS, pH 7.4 or 1% BSA/50% glycerol/PBS, pH 7.4).
- Mix by gently swirling the vial until content is completely dissolved.
- Adding 1ml of buffer will result in a Der f 2 concentration of 10,000ng/ml.



References:

1. Heyman PW, Chapman MD, Fox J, Aalberse, RC, PlattsMills, TAE. Antigenic and structural analysis of the Group II allergens (Derp II and Der f II) from house dust mites (*Dermatophagoides* sp). *J Allergy Clin Immunology* 1989;83:1055-67.
2. Mueller GA, Smith AM, Williams DC Jr., Hakkaart GAJ, Aalberse RC, Chapman MD, Rule, GS, Benjamin DC. Expression and secondary structure determination by NMR methods of the major house dust mite allergen Der p 2. *J Biol Chem* 1997;272:26893-8.
3. Smith AM et al. The molecular basis of antigenic cross-reactivity between the group 2 mite allergens. *J Allergy Clin Immunol* 2001;107:977-84.
4. Chapman MD et al. The European Union CREATE project: a model for international standardization of allergy diagnostics and vaccines. *J Allergy Clin Immunol.* 2008;122:882-889.
5. van Ree R et al. The CREATE project: development of certified reference materials for allergenic products and validation of methods for their quantification. *Allergy.* 2008;63(3):310-26.
6. Kaul S et al. Regulatory environment for allergen-specific immunotherapy. *Allergy* 2011;66:753-64.
7. Chapman MD and Briza P. Molecular approaches to allergen standardization. *Curr Allergy Asthma Rep.* 2012;12:478-84.
8. Chapman MD et al. Technological Innovations for High-Throughput Approaches to In Vitro Allergy Diagnosis. *Curr Allergy Asthma Rep.* 2015;15:36.